

Table 1C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa
Summary of Demographic Indicators ⁽²⁾

Page: 1
Run Date: 8 / 18 / 10

Total Records	Number	%	Source of Data	Number	%
Complete Records	14,992	73.4	WIC	20,415	100.0
Prenatal Records	1,511	7.4	Maternal Health	0	0.0
Postpartum Records	3,912	19.2	Other	0	0.0
Total Records Accepted	20,415		Total	20,415	100.0
			<i>Excluded Unknown: 0</i>	<i>0.0</i>	

Race/Ethnic Distribution	Number	%	Age Distribution	Number	%	Education Distribution	Number	%
White, Not Hispanic	14,622	71.9	< 15 Years	62	0.3	< High School	5,629	29.4
Black, Not Hispanic	1,630	8.0	15 - 17 Years	1,238	6.1	High School	8,446	44.1
Hispanic	3,331	16.4	18 - 19 Years	2,800	13.7	> High School	5,091	26.6
American Indian/Alaska Native	103	0.5	20 - 29 Years	12,701	62.2	Total	19,166	100.0
Asian/Pacific Islander	391	1.9	30 - 39 Years	3,400	16.7			
Multiple Races	259	1.3	>= 40 Years	209	1.0			
All Other	0	0.0	Total	20,410	100.0			
Total	20,336	100.0	<i>Excluded Unknown: 5</i>	<i>0.0</i>		<i>Excluded Unknown: 1,249</i>	<i>6.1</i>	
<i>Excluded Unknown: 79</i>	<i>0.4</i>							

% Poverty Level	Number	%	Program Participation ⁽³⁾	Number	%	Migrant Status	Number	%
0 - 50	5,978	36.4	WIC	16,503	100.0	Yes	4	0.0
51 - 100	5,077	30.9	SNAP	16,480	25.6	No	20,373	100.0
101 - 130	2,287	13.9	Medicaid	16,480	67.3	Total	20,377	100.0
131 - 150	1,176	7.2	TANF	16,480	4.5			
151 - 185	1,417	8.6	<i>Excluded Unknown:</i>					
186 - 200	162	1.0	WIC:	0	0.0			
> 200	346	2.1	SNAP:	23	0.1			
Adjunctive Eligibility	0	0.0	Medicaid:	23	0.1			
Total	16,443	100.0	TANF:	23	0.1	<i>Excluded Unknown: 38</i>	<i>0.2</i>	
<i>Excluded Unknown: 60</i>	<i>0.4</i>							

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Prenatal program participation.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 2C
2009 Pregnancy Nutrition Surveillance (1)
Iowa
Summary of Health Indicators

Summary of Health Indicators (2)				
	Iowa Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Prepregnancy BMI (3)				
Underweight	17,397	2.6	1,224,766	4.7
Normal Weight	17,397	39.2	1,224,766	43.1
Overweight	17,397	25.4	1,224,766	25.7
Obese	17,397	32.8	1,224,766	26.4
Maternal Weight Gain (4)				
Less Than Ideal	15,582	16.7	963,624	21.4
Ideal	15,582	31.3	963,624	30.3
Greater Than Ideal	15,582	52.1	963,624	48.2
Anemia (Low Hb/Hct) (5)				
1st Trimester	6,636	4.9	394,095	7.6
2nd Trimester	5,776	9.8	354,690	12.1
3rd Trimester	3,275	30.1	188,530	33.8
Postpartum	6,176	22.9	416,710	29.6
Parity (6)				
0	15,557	37.5	1,149,553	42.6
1	15,557	28.0	1,149,553	26.4
2	15,557	18.3	1,149,553	16.0
≥ 3	15,557	16.2	1,149,553	14.9
Interpregnancy Interval				
< 6 Months	11,824	15.7	618,788	12.6
6 - <18 Months	11,824	29.3	618,788	27.1
≥18 Months	11,824	55.0	618,788	60.3
Gestational Diabetes	17,786	7.1	346,704	4.4
Hypertension During Pregnancy	17,792	11.8	395,960	5.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

(6) Number of times a woman has given birth to an infant 20 or more weeks gestation.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 2C
2009 Pregnancy Nutrition Surveillance (1)
Iowa
Summary of Health Indicators

Summary of Health Indicators (2)				
	Iowa Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Medical Care				
1st Trimester	12,363	88.1	580,485	80.1
No Care	12,363	0.0	580,485	3.1
WIC Enrollment				
1st Trimester	17,948	44.1	1,187,387	31.7
2nd Trimester	17,948	30.5	1,187,387	34.0
3rd Trimester	17,948	16.4	1,187,387	18.4
Postpartum (3)	17,948	9.0	1,187,387	16.0
Prepregnancy Multivitamin Consumption (4)	14,555	15.9	266,567	19.3
Multivitamin Consumption During Pregnancy (5)	13,273	6.2	269,548	80.6
Smoking				
3 Months Prior to Pregnancy	15,549	39.8	999,199	23.1
During Pregnancy	15,280	25.5	926,915	14.2
Last 3 Months of Pregnancy	17,863	23.1	856,953	14.2
Postpartum	17,973	24.7	839,512	16.1
Smoking Changes (6)				
Quit Smoking by First Prenatal Visit	5,920	36.4	193,938	42.1
Quit Smoking by First Prenatal Visit and Stayed Off Cigarettes	5,069	30.7	146,933	35.5
Smoking in Household				
During Pregnancy	15,142	20.8	655,095	18.9
Postpartum	17,694	8.8	578,852	19.9
Drinking				
3 Months Prior to Pregnancy	9,008	14.8	975,737	7.8
Last 3 Months of Pregnancy	17,789	0.4	701,124	0.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Women who initially enrolled in WIC during the postpartum period.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

(6) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 2C
2009 Pregnancy Nutrition Surveillance (1)
Iowa
Summary of Health Indicators

Page: 4
Run Date: 8 / 18 / 10

Summary of Health Indicators (2)				
	Iowa Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Birthweight (3)				
Very Low	13,440	0.9	1,085,134	1.1
Low	13,440	6.8	1,085,134	7.1
Normal	13,440	83.9	1,085,134	85.2
High	13,440	8.5	1,085,134	6.6
Preterm (4)	12,069	10.2	1,025,278	11.5
Full term Low Birth Weight (5)	10,392	3.2	890,981	3.6
Multiple Births	18,904	1.1	1,060,021	1.4
Breastfeeding				
Ever Breastfed	14,299	33.3	952,980	65.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Very low birth weight (<1500 g), low birth weight (1500 - <2500 g), normal birth weight (2500 - <4000 g), high birth weight (>4000 g).

(4) <37 weeks gestation.

(5) 37 or more weeks gestation and <2500 g.



Table 3C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa

Page: 5
Run Date: 8 / 18 / 10

Comparison of Racial and Ethnic Distribution by Contributor to Nation

Comparison of Racial and Ethnic Distribution ⁽²⁾								
	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Iowa	20,415	71.6	8.0	16.3	0.5	1.9	1.3	0.4
Nation (Prior Year)	1,307,887	42.0	24.2	28.1	1.0	3.1	1.1	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa
Comparison of Age Distribution by Contributor to Nation

Comparison of Age Distribution ⁽²⁾								
	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Iowa	20,415	0.3	6.1	13.7	62.2	16.7	1.0	0.0
Nation (Prior Year)	1,307,887	0.3	6.3	12.6	59.7	19.4	1.4	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa

Page: 7
Run Date: 8 / 18 / 10

Comparison of Maternal Health Indicators by Contributor to Nation

Comparison of Maternal Health Indicators ⁽²⁾														
	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
Iowa	17,397	2.6	58.2	15,582	16.7	52.1	3,275	30.1	6,176	22.9	17,786	7.1	17,792	11.8
Nation (Prior Year)	1,224,766	4.7	52.1	963,624	21.4	48.2	188,530	33.8	416,710	29.6	346,704	4.4	395,960	5.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa

Page: 8
Run Date: 8 / 18 / 10

Comparison of Maternal Behavioral Indicators by Contributor to Nation

Comparison of Maternal Behavioral Indicators ⁽²⁾								
	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (3)	During Pregnancy (4)	
	Number	%	%	Number	%	Number	%	Number
Iowa	12,363	88.1	0.0	17,948	44.1	14,555	15.9	13,273
Nation (Prior Year)	580,485	80.1	3.1	1,187,387	31.7	266,567	19.3	269,548

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(4) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7C
2009 Pregnancy Nutrition Surveillance (1)
Iowa
Comparison of Smoking Indicators by Contributor to Nation

Comparison of Smoking Indicators (2)										
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	%	Number	%	Number	%	Number	%	Number	%
Iowa	15,549	39.8	17,863	23.1	5,069	30.7	15,142	20.8	17,694	8.8
Nation (Prior Year)	999,199	23.1	856,953	14.2	146,933	35.5	655,095	18.9	578,852	19.9

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾

Page: 10
Run Date: 8 / 18 / 10

Iowa
Comparison of Infant Health Indicators by Contributor to Nation

Comparison of Infant Health Indicators ⁽²⁾									
	Birthweight			Preterm (5)	Full Term LBW (6)	Ever Breastfed			
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
Iowa	13,440	7.6	8.5	12,069	10.2	10,392	3.2	14,299	33.3
Nation (Prior Year)	1,085,134	8.2	6.6	1,025,278	11.5	890,981	3.6	952,980	65.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 9C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa

Maternal Health Indicators by Race/Ethnicity, Age, or Education

Maternal Health Indicators by Race/Ethnicity, Age, or Education ⁽²⁾

	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)				Gestational Diabetes		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
Race/Ethnicity														
White, Not Hispanic	12,403	2.9	58.0	11,172	15.5	54.5	2,336	27.7	4,522	19.9	12,914	6.8	12,918	12.9
Black, Not Hispanic	1,415	2.5	58.7	1,230	21.0	48.9	264	56.8	440	45.9	1,366	5.2	1,367	11.4
Hispanic	2,971	1.0	61.3	2,651	19.6	43.8	539	27.6	961	24.3	2,870	8.9	2,870	8.0
American Indian/Alaska Native	88	*	*	71	*	*	20	*	47	*	82	*	83	*
Asian/Pacific Islander	290	4.1	31.4	261	21.5	39.8	82	*	138	24.6	332	8.1	332	4.8
Multiple Races	224	2.7	57.6	197	16.2	59.4	34	*	68	*	222	5.9	222	7.7
All Other/Unknown	6	*	*	0	*	*	0	*	0	*	0	*	0	*
Total	17,397	2.6	58.2	15,582	16.7	52.1	3,275	30.1	6,176	22.9	17,786	7.1	17,792	11.8
Age														
< 15 Years	50	*	*	41	*	*	11	*	10	*	49	*	49	*
15 - 17 Years	1,080	4.5	37.5	986	15.0	57.9	189	36.5	330	35.5	1,071	2.6	1,071	12.6
18 - 19 Years	2,497	4.2	46.3	2,239	15.5	55.1	461	32.3	742	29.4	2,406	3.0	2,408	11.2
20 - 29 Years	10,775	2.4	60.2	9,637	16.5	52.0	2,079	30.0	3,902	22.1	11,113	6.3	11,115	11.6
30 - 39 Years	2,832	1.1	68.9	2,539	18.9	47.5	507	26.0	1,132	17.1	2,965	14.2	2,967	12.1
>= 40 Years	158	0.0	69.6	136	20.6	44.9	27	*	60	*	178	18.0	178	21.9
Unknown	5	*	*	4	*	*	1	*	0	*	4	*	4	*
Total	17,397	2.6	58.2	15,582	16.7	52.1	3,275	30.1	6,176	22.9	17,786	7.1	17,792	11.8
Education														
< High School	4,994	3.3	53.8	4,485	18.3	49.6	896	35.4	1,589	27.9	4,881	6.1	4,883	10.1
High School	7,377	2.4	60.2	6,635	15.9	53.2	1,397	29.6	2,423	23.6	7,430	7.4	7,436	12.3
> High School	4,115	2.3	60.5	3,681	15.8	53.5	825	24.1	1,783	17.6	4,479	7.3	4,478	13.3
Unknown	911	2.0	56.1	781	17.9	49.8	157	35.7	381	21.3	996	7.5	995	9.6
Total	17,397	2.6	58.2	15,582	16.7	52.1	3,275	30.1	6,176	22.9	17,786	7.1	17,792	11.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for health indicators, and data with errors for health and demographic indicators.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 10C
2009 Pregnancy Nutrition Surveillance (1)
Iowa

Maternal Behavioral Indicators by Race/Ethnicity, Age, or Education

Maternal Behavioral Indicators by Race/Ethnicity, Age, or Education (2)									
	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (3)		During Pregnancy (4)	
	Number	%	%	Number	%	Number	%	Number	%
Race/Ethnicity									
White, Not Hispanic	8,966	90.2	0.0	12,741	45.2	10,390	17.6	9,623	7.1
Black, Not Hispanic	944	82.2	0.0	1,450	34.6	1,147	9.0	982	4.4
Hispanic	2,029	83.8	0.0	3,021	44.5	2,498	12.2	2,188	3.5
American Indian/Alaska Native	56	*	*	92	*	70	*	69	*
Asian/Pacific Islander	222	78.8	0.0	331	33.2	266	18.0	236	3.0
Multiple Races	146	82.2	0.0	234	47.0	183	15.3	174	6.3
All Other/Unknown	0	*	*	79	*	1	*	1	*
Total	12,363	88.1	00.0	17,948	44.1	14,555	15.9	13,273	6.2
Age									
< 15 Years	32	*	*	58	*	42	*	42	*
15 - 17 Years	802	81.4	0.0	1,134	46.9	918	6.0	848	5.8
18 - 19 Years	1,793	87.1	0.0	2,590	47.9	2,119	8.4	2,002	6.0
20 - 29 Years	7,703	88.9	0.0	11,095	43.0	9,070	16.2	8,147	6.5
30 - 39 Years	1,932	89.3	0.0	2,894	43.8	2,273	25.6	2,117	5.6
>= 40 Years	98	*	*	172	42.4	129	24.0	113	3.5
Unknown	3	*	*	5	*	4	*	4	*
Total	12,363	88.1	00.0	17,948	44.1	14,555	15.9	13,273	6.2
Education									
< High School	3,562	84.3	0.0	5,136	45.6	4,267	9.5	3,693	5.7
High School	5,408	88.9	0.0	7,570	45.4	6,278	14.7	5,758	6.3
> High School	2,890	91.2	0.0	4,247	41.3	3,368	26.9	3,230	6.2
Unknown	503	88.5	0.0	995	38.0	642	12.5	592	7.8
Total	12,363	88.1	00.0	17,948	44.1	14,555	15.9	13,273	6.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for behavioral indicators, and data with errors for behavioral and demographic indicators.

(3) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(4) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 11C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa
Smoking Indicators by Race/Ethnicity, Age, or Education

Smoking Indicators by Race/Ethnicity, Age, or Education ⁽²⁾										
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	%	Number	%		Number	%	Number	%	
Race/Ethnicity										
White, Not Hispanic	10,994	48.2	12,992	27.9	4,396	29.3	10,690	24.4	12,869	10.4
Black, Not Hispanic	1,231	29.7	1,369	16.3	268	36.6	1,198	15.2	1,365	5.6
Hispanic	2,762	13.4	2,871	6.6	291	43.3	2,714	9.3	2,836	3.8
American Indian/Alaska Native	82	*	83	*	28	*	79	*	81	*
Asian/Pacific Islander	286	10.5	328	5.5	21	*	275	12.0	325	4.3
Multiple Races	193	45.1	220	27.3	65	*	185	22.2	218	9.6
All Other/Unknown	1	*	0	*	0	*	1	*	0	*
Total	15,549	39.8	17,863	23.1	5,069	30.7	15,142	20.8	17,694	8.8
Age										
< 15 Years	48	*	49	*	5	*	49	*	48	*
15 - 17 Years	1,004	34.3	1,072	17.3	299	41.8	993	38.5	1,071	15.7
18 - 19 Years	2,265	45.5	2,422	23.5	868	35.7	2,257	29.6	2,402	10.9
20 - 29 Years	9,666	41.9	11,168	24.5	3,288	29.5	9,333	18.4	11,043	8.5
30 - 39 Years	2,427	30.1	2,967	20.2	582	23.9	2,375	14.0	2,947	5.7
>= 40 Years	135	23.7	181	14.4	26	*	131	16.0	179	7.8
Unknown	4	*	4	*	1	*	4	*	4	*
Total	15,549	39.8	17,863	23.1	5,069	30.7	15,142	20.8	17,694	8.8
Education										
< High School	4,673	39.5	4,900	25.6	1,509	25.8	4,534	24.9	4,855	11.1
High School	6,660	44.8	7,456	26.5	2,464	30.4	6,457	22.1	7,396	9.5
> High School	3,514	31.7	4,505	15.1	911	40.0	3,453	12.6	4,454	5.5
Unknown	702	35.8	1,002	21.5	185	29.2	698	21.6	989	7.6
Total	15,549	39.8	17,863	23.1	5,069	30.7	15,142	20.8	17,694	8.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for smoking indicators, and data with errors for smoking and demographic indicators.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 12C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa
Infant Health Indicators by Race/Ethnicity, Age, or Education

Infant Health Indicators by Race/Ethnicity, Age, or Education ⁽²⁾									
	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)	High (4)							
	Number	%	%	Number	%	Number	%	Number	%
Race/Ethnicity									
White, Not Hispanic	9,842	7.5	8.8	8,767	10.2	7,562	2.9	10,455	30.4
Black, Not Hispanic	966	11.9	4.9	884	13.2	727	6.6	1,041	28.0
Hispanic	2,144	6.4	9.2	1,993	9.5	1,728	3.1	2,290	48.3
American Indian/Alaska Native	66	*	*	59	*	52	*	71	*
Asian/Pacific Islander	259	6.2	2.7	214	8.4	192	2.6	269	44.2
Multiple Races	163	8.6	7.4	152	8.6	131	6.1	173	27.7
All Other/Unknown	0	*	*	0	*	0	*	0	*
Total	13,440	7.6	8.5	12,069	10.2	10,392	3.2	14,299	33.3
Age									
< 15 Years	41	*	*	40	*	35	*	42	*
15 - 17 Years	782	9.0	5.1	732	10.7	629	4.5	821	24.1
18 - 19 Years	1,784	6.7	6.6	1,724	10.0	1,476	3.0	1,915	27.2
20 - 29 Years	8,467	6.5	8.5	7,545	9.8	6,534	3.2	9,007	33.0
30 - 39 Years	2,245	7.2	11.0	1,925	11.7	1,635	3.1	2,374	42.4
>= 40 Years	117	6.0	12.0	99	*	79	*	136	36.0
Unknown	4	*	*	4	*	4	*	4	*
Total	13,440	7.6	8.5	12,069	10.2	10,392	3.2	14,299	33.3
Education									
< High School	3,643	7.2	6.5	3,432	10.9	2,929	3.5	3,914	29.4
High School	5,672	6.8	8.8	5,209	9.5	4,525	3.5	6,031	29.0
> High School	3,387	6.1	10.1	2,855	10.3	2,463	2.4	3,569	44.1
Unknown	738	7.6	8.7	573	12.7	475	2.9	785	36.4
Total	13,440	7.6	8.5	12,069	10.2	10,392	3.2	14,299	33.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for health indicators, and data with errors for health and demographic indicators.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 13C
2009 Pregnancy Nutrition Surveillance (1)
Iowa

Maternal Weight Gain and Birth Outcomes by Select Health Indicators

Maternal Weight Gain and Birth Outcomes by Select Health Indicators ⁽²⁾										
	Weight Gain (3)			Birth Outcomes						
	< Ideal		> Ideal	Birthweight			Preterm (6)		Full Term LBW (7)	
				Low (4)	High (5)					
		Number	%	%	Number	%	%	Number	%	Number
Prepregnancy BMI										
Underweight	401	25.2	30.4	292	15.1	3.4	303	12.5	249	6.8
Normal Weight	6,178	20.8	41.4	4,412	8.0	5.9	4,559	10.7	3,889	3.7
Overweight	4,001	9.5	62.4	2,880	5.8	9.1	2,949	9.0	2,593	2.2
Obese	5,002	16.7	58.7	3,739	6.5	11.9	3,828	9.9	3,313	2.7
Weight Gain										
< Ideal	–	–	–	1,694	13.3	4.0	1,760	15.9	1,411	6.0
Ideal	–	–	–	3,513	8.3	5.5	3,631	10.5	3,111	3.7
> Ideal	–	–	–	5,945	4.7	11.9	6,073	8.1	5,367	1.9
Smoking During Pregnancy										
Smokers	3,300	18.1	51.5	2,465	9.2	4.3	2,530	10.7	2,174	5.2
Non-Smokers	9,946	16.3	52.1	7,332	6.1	10.0	7,570	9.4	6,571	2.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 14C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa
Summary of Trends in Racial and Ethnic Distribution

Page: 16
Run Date: 8 / 18 / 10

Summary of Trends in Racial and Ethnic Distribution ⁽²⁾								
Year	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
2009	20,415	71.6	8.0	16.3	0.5	1.9	1.3	0.4
2008	20,660	71.9	7.4	17.0	0.5	1.8	1.2	0.3
2007	20,774	71.8	7.5	17.1	0.5	1.5	1.2	0.4
2006	11,150	71.9	6.8	17.9	0.3	1.8	1.1	0.2
2005	17,919	75.9	7.3	14.3	0.7	1.8	0.0	0.0
2004	13,126	76.3	7.2	13.5	0.7	2.2	0.0	0.0
2003	18,177	78.0	6.8	12.2	0.6	2.4	0.0	0.0
2002	18,053	78.1	6.5	12.3	0.7	2.3	0.0	0.0
2001	16,916	79.7	6.8	11.0	0.6	1.9	0.0	0.0
2000	17,515	80.4	6.5	10.3	0.7	2.2	0.0	0.0
1999	17,411	81.7	6.2	9.5	0.6	2.1	0.0	0.0
1998	17,784	83.1	5.9	8.5	0.6	2.1	0.0	0.0
1997	17,515	82.8	6.1	8.2	0.7	2.3	0.0	0.0
1996	15,733	83.9	6.1	7.5	0.5	2.1	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 15C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa
Summary of Trends in Age Distribution

Summary of Trends in Age Distribution ⁽²⁾								
Year	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2009	20,415	0.3	6.1	13.7	62.2	16.7	1.0	0.0
2008	20,660	0.3	6.5	13.8	62.1	16.2	1.1	0.0
2007	20,774	0.3	5.8	13.8	63.2	15.8	1.0	0.1
2006	11,150	0.3	6.2	13.0	63.9	15.6	1.0	0.1
2005	17,919	0.3	6.5	13.5	63.2	15.5	1.0	0.0
2004	13,126	0.2	5.6	13.6	63.4	16.0	1.1	0.0
2003	18,177	0.3	6.6	14.2	62.8	15.3	0.8	0.0
2002	18,053	0.3	6.8	14.4	62.3	15.3	1.0	0.0
2001	16,916	0.3	7.0	15.5	61.4	14.9	0.8	0.0
2000	17,515	0.4	7.7	15.9	60.6	14.5	0.9	0.0
1999	17,411	0.4	8.0	16.3	59.8	14.7	0.8	0.0
1998	17,784	0.4	8.3	15.7	59.8	14.9	0.8	0.0
1997	17,515	0.5	8.9	15.5	59.6	14.8	0.8	0.0
1996	15,733	0.4	8.7	15.8	59.3	15.1	0.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 16C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa
Summary of Trends in Maternal Health Indicators

Summary of Trends in Maternal Health Indicators ⁽²⁾										
Year	Pregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
2009	17,397	2.6	58.2	15,582	16.7	52.1	3,275	30.1	6,176	22.9
2008	17,526	2.9	57.5	15,701	15.9	53.0	3,252	28.0	6,741	21.9
2007	17,366	2.7	56.2	15,268	16.1	51.8	3,057	33.0	6,908	24.2
2006	9,130	3.1	55.0	7,746	17.0	51.8	1,743	28.8	3,824	21.8
2005	16,560	5.2	52.2	12,705	17.9	51.4	2,655	30.2	5,986	25.3
2004	11,846	5.3	50.9	9,831	18.9	50.2	1,626	29.9	5,746	24.3
2003	16,572	5.4	50.2	14,498	17.9	50.6	2,496	26.8	7,630	23.5
2002	16,489	5.9	49.4	14,267	17.7	50.6	2,531	25.9	7,365	21.7
2001	15,867	6.4	49.4	13,502	18.6	50.0	2,473	25.2	6,394	21.7
2000	16,502	6.7	47.9	14,345	18.8	49.4	2,507	23.8	6,425	21.3
1999	16,328	6.7	46.5	14,246	18.9	48.8	2,358	24.2	5,732	19.6
1998	16,476	6.7	45.9	14,708	18.7	49.0	2,349	25.0	5,698	20.3
1997	16,031	6.5	44.7	14,318	18.7	47.1	2,182	22.0	5,832	19.9
1996	10,991	7.3	42.6	9,937	19.4	47.2	1,994	24.9	5,609	19.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 17C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa
Summary of Trends in Maternal Behavioral Indicators

Summary of Trends in Maternal Behavioral Indicators ⁽²⁾					
Year	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	%	%	Number	%
2009	12,363	88.1	0.0	17,948	44.1
2008	12,046	87.9	0.0	17,899	43.4
2007	11,812	86.5	0.0	15,981	41.7
2006	3,998	86.7	0.0	8,328	38.8
2005	14,170	81.4	11.0	17,429	35.4
2004	11,356	79.6	13.5	12,403	32.1
2003	16,442	79.9	12.7	17,405	33.9
2002	16,189	79.3	13.3	17,216	33.4
2001	14,906	80.3	11.6	16,326	32.8
2000	15,656	79.0	12.3	16,869	33.9
1999	15,514	78.8	12.5	16,691	34.2
1998	15,902	78.6	12.5	16,983	34.8
1997	15,523	78.1	13.0	16,687	34.7
1996	11,363	76.6	14.0	13,211	29.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 18C
2009 Pregnancy Nutrition Surveillance (1)
Iowa
Summary of Trends in Smoking Indicators

Page: 20
Run Date: 8 / 18 / 10

Summary of Trends in Smoking Indicators (2)										
Year	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	%	Number	%	Number	%	Number	%	Number	%
2009	15,549	39.8	17,863	23.1	5,069	30.7	15,142	20.8	17,694	8.8
2008	15,279	40.8	17,175	25.1	5,054	29.8	15,073	24.4	17,058	15.4
2007	15,104	42.8	18,082	25.6	4,619	30.7	12,455	26.3	17,812	16.8
2006	4,580	41.0	6,096	25.7	0	*	6	*	6,002	16.5
2005	17,506	42.1	14,294	27.1	4,827	29.4	14,443	41.9	14,289	36.5
2004	12,492	41.9	11,435	26.9	3,252	29.1	9,111	41.9	11,432	36.0
2003	17,451	43.1	16,563	27.1	5,542	29.9	13,966	40.8	16,550	35.8
2002	17,302	44.2	16,331	27.7	5,615	29.9	13,723	41.7	16,309	34.7
2001	16,419	46.1	15,026	29.8	5,393	27.3	13,049	44.4	15,021	36.4
2000	16,981	46.0	15,747	29.6	5,602	27.9	13,443	44.6	15,739	37.8
1999	16,817	46.1	15,630	29.9	5,670	28.9	13,414	47.4	15,618	39.7
1998	17,066	44.4	16,175	28.9	5,735	28.1	13,648	47.8	16,159	39.9
1997	16,574	42.9	15,888	28.9	5,372	26.9	13,216	47.6	15,884	39.9
1996	11,437	41.8	12,593	30.3	3,313	23.8	8,212	48.6	12,575	42.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 19C
2009 Pregnancy Nutrition Surveillance (1)
Iowa
Summary of Trends in Infant Health Indicators

Summary of Trends in Infant Health Indicators ⁽²⁾									
Year	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)	High (4)							
	Number	%	%	Number	%	Number	%	Number	%
2009	13,440	7.6	8.5	12,069	10.2	10,392	3.2	14,299	33.3
2008	13,645	7.9	8.0	11,797	11.1	10,307	3.0	14,114	29.7
2007	14,468	7.8	8.4	10,848	10.8	9,521	2.8	14,889	22.5
2006	4,848	7.2	8.4	3,502	10.1	3,098	2.5	4,968	25.6
2005	13,861	7.4	8.3	13,462	13.1	11,670	3.1	13,858	58.1
2004	11,041	7.8	8.2	10,402	13.0	9,025	2.9	11,030	58.6
2003	16,030	6.9	9.0	15,405	12.4	13,445	2.7	16,048	57.7
2002	15,796	7.3	9.2	15,112	12.8	13,115	2.9	15,800	57.2
2001	14,604	6.9	9.5	14,082	12.3	12,327	3.1	14,577	55.7
2000	15,264	6.9	9.6	14,708	11.5	12,983	2.8	15,245	54.1
1999	15,183	6.9	10.1	14,548	11.7	12,813	2.6	15,159	52.5
1998	15,680	6.9	10.6	14,966	11.8	13,180	2.9	15,624	52.4
1997	15,378	6.6	10.1	14,657	11.7	12,925	2.8	15,356	49.8
1996	12,243	6.8	10.1	11,477	11.3	10,169	3.1	12,197	32.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa

Page: 22
Run Date: 8 / 18 / 10

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)		
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum
		Number	%	%	Number	%	%	Number	%	Number %
White, Not Hispanic	2009	12,403	2.9	58.0	11,172	15.5	54.5	2,336	27.7	4,522 19.9
	2008	12,532	3.2	57.3	11,203	14.6	55.7	2,299	26.5	4,831 19.5
	2007	12,463	3.1	55.9	10,957	15.2	54.1	2,171	30.8	4,863 21.3
	2006	6,552	3.5	54.5	5,564	15.7	54.6	1,252	24.8	2,715 19.8
	2005	12,732	5.5	52.3	9,999	16.7	53.6	1,964	27.2	4,500 23.5
	2004	9,207	5.5	50.9	7,781	17.6	52.4	1,197	27.8	4,310 22.0
	2003	13,096	5.8	50.0	11,626	16.8	52.3	1,877	25.3	5,833 22.1
	2002	13,052	6.2	49.3	11,460	16.7	52.6	1,946	24.5	5,602 20.1
	2001	12,746	6.8	49.6	10,999	17.6	51.7	1,935	23.6	5,008 19.7
	2000	13,363	7.1	47.4	11,800	17.8	50.9	1,922	22.2	5,083 19.1
Black, Not Hispanic	2009	1,415	2.5	58.7	1,230	21.0	48.9	264	56.8	440 45.9
	2008	1,311	4.0	58.7	1,174	19.8	51.7	248	49.2	451 41.7
	2007	1,310	2.2	57.6	1,132	17.7	50.7	268	56.0	506 47.8
	2006	618	3.4	56.3	524	19.7	48.1	133	49.6	236 37.3
	2005	1,203	4.9	56.4	877	20.4	46.4	226	47.8	398 41.5
	2004	818	4.6	54.3	645	22.0	46.8	146	44.5	376 42.6
	2003	1,094	3.7	53.9	933	19.3	50.7	203	48.8	473 42.5
	2002	1,043	4.3	54.2	883	17.2	52.8	166	49.4	444 40.3
	2001	1,073	5.8	52.4	869	21.2	49.5	171	45.0	374 36.9
	2000	1,053	5.2	53.8	881	22.1	49.7	171	40.9	378 39.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa

Page: 23
Run Date: 8 / 18 / 10

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)		
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum
		Number	%	%	Number	%	%	Number	%	Number %
Hispanic	2009	2,971	1.0	61.3	2,651	19.6	43.8	539	27.6	961 24.3
	2008	3,115	1.2	60.7	2,821	19.9	43.4	555	25.4	1,172 23.2
	2007	3,076	1.4	58.7	2,735	18.8	44.1	491	30.3	1,292 25.5
	2006	1,690	1.1	58.4	1,438	21.5	43.3	298	33.2	723 23.4
	2005	2,223	3.1	53.8	1,535	23.7	43.2	391	33.2	934 27.0
	2004	1,477	2.6	54.6	1,134	24.0	40.7	221	28.5	872 28.6
	2003	1,887	2.8	54.5	1,512	23.4	42.9	322	24.5	1,032 20.9
	2002	1,907	3.3	53.0	1,523	22.8	39.9	333	24.0	1,040 22.0
	2001	1,656	2.7	50.9	1,308	22.1	41.4	299	22.4	811 25.0
	2000	1,632	3.1	53.2	1,291	23.4	40.4	322	23.0	759 24.9
American Indian/Alaska Native	2009	88	*	*	71	*	*	20	*	47 *
	2008	95	*	*	82	*	*	20	*	50 *
	2007	90	*	*	69	*	*	15	*	37 *
	2006	27	*	*	24	*	*	6	*	16 *
	2005	105	1.0	49.5	75	*	*	20	*	43 *
	2004	85	*	*	58	*	*	21	*	40 *
	2003	99	*	*	72	*	*	14	*	57 *
	2002	108	6.5	64.8	79	*	*	19	*	60 *
	2001	90	*	*	62	*	*	17	*	42 *
	2000	102	2.9	53.9	59	*	*	17	*	44 *

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa

Page: 24
Run Date: 8 / 18 / 10

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)		
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum
		Number	%	%	Number	%	%	Number	%	Number %
Asian/Pacific Islander	2009	290	4.1	31.4	261	21.5	39.8	82	*	138 24.6
	2008	250	3.2	24.4	231	16.9	39.8	84	*	163 23.9
	2007	213	5.2	25.4	182	18.7	34.1	61	*	124 20.2
	2006	133	4.5	35.3	107	17.8	41.1	33	*	89 *
	2005	297	12.8	19.5	219	25.1	30.1	54	*	111 20.7
	2004	259	11.2	19.7	213	32.9	27.7	41	*	148 19.6
	2003	396	10.6	21.2	355	25.1	29.0	80	*	235 24.7
	2002	379	12.7	16.1	322	31.4	26.1	67	*	219 18.3
	2001	302	13.6	18.9	264	31.8	26.9	51	*	159 25.2
	2000	352	14.5	20.2	314	26.8	28.3	75	*	161 27.3
Multiple Races	2009	224	2.7	57.6	197	16.2	59.4	34	*	68 *
	2008	220	2.7	53.2	190	10.5	60.5	46	*	73 *
	2007	209	3.3	55.5	193	17.6	50.8	51	*	86 *
	2006	101	4.0	50.5	88	*	*	20	*	44 *
	2005	0	*	*	0	*	*	0	*	0 *
	2004	0	*	*	0	*	*	0	*	0 *
	2003	0	*	*	0	*	*	0	*	0 *
	2002	0	*	*	0	*	*	0	*	0 *
	2001	0	*	*	0	*	*	0	*	0 *
	2000	0	*	*	0	*	*	0	*	0 *

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa

Page: 25
Run Date: 8 / 18 / 10

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
All Other/Unknown	2009	6	*	*	0	*	*	0	*	0	*
	2008	3	*	*	0	*	*	0	*	1	*
	2007	5	*	*	0	*	*	0	*	0	*
	2006	9	*	*	1	*	*	1	*	1	*
	2005	0	*	*	0	*	*	0	*	0	*
	2004	0	*	*	0	*	*	0	*	0	*
	2003	0	*	*	0	*	*	0	*	0	*
	2002	0	*	*	0	*	*	0	*	0	*
	2001	0	*	*	0	*	*	0	*	0	*
	2000	0	*	*	0	*	*	0	*	0	*
Total All Race/Ethnic Groups	2009	17,397	2.6	58.2	15,582	16.7	52.1	3,275	30.1	6,176	22.9
	2008	17,526	2.9	57.5	15,701	15.9	53.0	3,252	28.0	6,741	21.9
	2007	17,366	2.7	56.2	15,268	16.1	51.8	3,057	33.0	6,908	24.2
	2006	9,130	3.1	55.0	7,746	17.0	51.8	1,743	28.8	3,824	21.8
	2005	16,560	5.2	52.2	12,705	17.9	51.4	2,655	30.2	5,986	25.3
	2004	11,846	5.3	50.9	9,831	18.9	50.2	1,626	29.9	5,746	24.3
	2003	16,572	5.4	50.2	14,498	17.9	50.6	2,496	26.8	7,630	23.5
	2002	16,489	5.9	49.4	14,267	17.7	50.6	2,531	25.9	7,365	21.7
	2001	15,867	6.4	49.4	13,502	18.6	50.0	2,473	25.2	6,394	21.7
	2000	16,502	6.7	47.9	14,345	18.8	49.4	2,507	23.8	6,425	21.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa

Page: 26
Run Date: 8 / 18 / 10

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
White, Not Hispanic	2009	8,966	90.2	0.0	12,741	45.2
	2008	8,689	89.7	0.0	12,744	44.4
	2007	8,598	88.3	0.0	11,323	43.3
	2006	2,835	88.6	0.0	5,908	40.3
	2005	10,907	82.7	10.7	13,229	36.7
	2004	8,724	81.2	13.0	9,514	32.8
	2003	12,866	80.9	12.5	13,604	34.9
	2002	12,693	80.5	13.0	13,473	34.5
	2001	11,947	81.0	11.8	13,008	33.8
	2000	12,656	80.0	12.5	13,571	35.1
Black, Not Hispanic	2009	944	82.2	0.0	1,450	34.6
	2008	868	83.3	0.0	1,336	37.9
	2007	852	80.0	0.0	1,182	33.8
	2006	261	79.3	0.0	538	29.6
	2005	994	76.8	11.7	1,277	28.9
	2004	798	74.9	13.5	880	25.9
	2003	1,109	78.9	10.5	1,186	27.3
	2002	1,056	77.1	11.6	1,117	27.5
	2001	974	80.8	7.9	1,122	29.9
	2000	993	77.6	9.7	1,097	29.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa

Page: 27
Run Date: 8 / 18 / 10

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Hispanic	2009	2,029	83.8	0.0	3,021	44.5
	2008	2,110	83.6	0.0	3,158	42.7
	2007	2,015	83.2	0.0	2,900	38.6
	2006	771	82.6	0.0	1,598	37.4
	2005	1,933	76.9	12.2	2,491	32.8
	2004	1,514	75.0	15.9	1,649	32.9
	2003	1,980	74.6	15.5	2,099	33.0
	2002	1,972	73.4	16.1	2,114	31.0
	2001	1,613	76.5	11.3	1,794	28.9
	2000	1,586	72.1	13.6	1,724	29.6
American Indian/Alaska Native	2009	56	*	*	92	*
	2008	60	*	*	92	*
	2007	46	*	*	81	*
	2006	11	*	*	27	*
	2005	90	*	*	120	29.2
	2004	67	*	*	88	*
	2003	86	*	*	103	33.0
	2002	98	*	*	117	28.2
	2001	71	*	*	95	*
	2000	73	*	*	113	37.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa

Page: 28
Run Date: 8 / 18 / 10

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Asian/Pacific Islander	2009	222	78.8	0.0	331	33.2
	2008	176	75.6	0.0	299	30.8
	2007	157	80.9	0.0	224	35.3
	2006	60	*	*	133	32.3
	2005	246	83.3	11.0	311	26.7
	2004	253	75.5	14.6	272	22.8
	2003	401	78.3	12.7	413	22.5
	2002	370	77.8	12.4	395	25.3
	2001	301	75.4	15.0	307	24.1
	2000	348	78.7	8.3	364	20.9
Multiple Races	2009	146	82.2	0.0	234	47.0
	2008	143	86.0	0.0	219	41.6
	2007	144	76.4	0.0	188	37.2
	2006	60	*	*	98	*
	2005	0	*	*	0	*
	2004	0	*	*	0	*
	2003	0	*	*	0	*
	2002	0	*	*	0	*
	2001	0	*	*	0	*
	2000	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Iowa

Page: 29
Run Date: 8 / 18 / 10

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
All Other/Unknown	2009	0	*	*	79	*
	2008	0	*	*	51	*
	2007	0	*	*	83	*
	2006	0	*	*	26	*
	2005	0	*	*	1	*
	2004	0	*	*	0	*
	2003	0	*	*	0	*
	2002	0	*	*	0	*
	2001	0	*	*	0	*
	2000	0	*	*	0	*
Total All Race/Ethnic Groups	2009	12,363	88.1	0.0	17,948	44.1
	2008	12,046	87.9	0.0	17,899	43.4
	2007	11,812	86.5	0.0	15,981	41.7
	2006	3,998	86.7	0.0	8,328	38.8
	2005	14,170	81.4	11.0	17,429	35.4
	2004	11,356	79.6	13.5	12,403	32.1
	2003	16,442	79.9	12.7	17,405	33.9
	2002	16,189	79.3	13.3	17,216	33.4
	2001	14,906	80.3	11.6	16,326	32.8
	2000	15,656	79.0	12.3	16,869	33.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 22C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾

Page: 30
Run Date: 8 / 18 / 10

Iowa
Summary of Trends in Smoking Indicators by Race/Ethnicity

Summary of Trends in Smoking Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
White, Not Hispanic	2009	10,994	48.2	12,992	27.9	4,396	29.3	10,690	24.4	12,869	10.4
	2008	10,834	49.4	12,607	30.1	4,393	28.1	10,703	28.8	12,517	17.6
	2007	10,709	52.0	13,126	30.7	3,987	29.5	8,881	30.4	12,944	19.3
	2006	3,129	52.0	4,341	31.9	0	*	2	*	4,290	19.2
	2005	13,288	50.4	10,989	32.7	4,427	28.2	10,881	48.4	10,985	42.2
	2004	9,572	49.3	8,780	32.2	2,974	27.9	6,916	47.7	8,778	41.3
	2003	13,643	49.9	12,947	32.1	5,046	28.6	10,818	46.5	12,940	40.7
	2002	13,547	51.6	12,797	32.9	5,192	28.7	10,712	46.9	12,782	39.7
	2001	13,089	53.0	12,041	34.9	4,982	26.0	10,345	49.6	12,038	41.0
	2000	13,673	52.1	12,733	33.9	5,132	27.0	10,735	49.3	12,727	42.1
Black, Not Hispanic	2009	1,231	29.7	1,369	16.3	268	36.6	1,198	15.2	1,365	5.6
	2008	1,118	29.7	1,222	18.5	243	38.7	1,088	18.1	1,227	11.5
	2007	1,133	32.9	1,328	21.3	254	31.5	918	20.5	1,301	12.1
	2006	296	28.7	417	17.7	0	*	1	*	405	12.8
	2005	1,285	33.3	1,002	19.7	255	34.9	1,041	34.7	1,002	27.8
	2004	895	36.3	800	22.0	175	32.6	632	32.9	800	27.4
	2003	1,184	37.8	1,116	22.5	307	30.9	941	31.0	1,115	26.3
	2002	1,121	35.4	1,057	20.9	254	36.6	860	32.0	1,055	23.9
	2001	1,127	38.3	982	20.3	276	39.5	894	33.2	982	23.6
	2000	1,096	37.9	997	24.2	289	28.7	888	34.9	996	23.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 22C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾

Page: 31
Run Date: 8 / 18 / 10

Iowa
Summary of Trends in Smoking Indicators by Race/Ethnicity

Summary of Trends in Smoking Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Hispanic	2009	2,762	13.4	2,871	6.6	291	43.3	2,714	9.3	2,836	3.8
	2008	2,812	14.6	2,751	7.6	306	45.8	2,772	10.8	2,730	7.1
	2007	2,795	14.0	3,059	7.2	286	46.9	2,277	12.7	3,010	8.2
	2006	994	12.3	1,138	6.9	0	*	3	*	1,115	8.6
	2005	2,500	6.8	1,964	3.3	102	57.8	2,179	15.6	1,963	11.7
	2004	1,662	7.0	1,529	2.8	63	*	1,328	16.8	1,528	12.8
	2003	2,105	8.1	2,007	2.7	127	63.8	1,813	15.0	2,002	13.1
	2002	2,121	8.3	1,997	3.3	126	59.5	1,776	18.1	1,992	12.4
	2001	1,800	8.2	1,627	3.0	100	51.0	1,513	18.4	1,625	13.0
	2000	1,734	10.3	1,592	4.3	131	55.7	1,457	20.7	1,591	16.3
American Indian/Alaska Native	2009	82	*	83	*	28	*	79	*	81	*
	2008	86	*	86	*	29	*	80	*	83	*
	2007	74	*	89	*	18	*	57	*	85	*
	2006	16	*	23	*	0	*	0	*	21	*
	2005	121	34.7	91	*	25	*	95	*	91	*
	2004	88	*	68	*	18	*	65	*	68	*
	2003	103	43.7	86	*	27	*	82	*	86	*
	2002	117	37.6	99	*	21	*	92	*	99	*
	2001	95	*	72	*	17	*	72	*	72	*
	2000	113	43.4	73	*	19	*	94	*	73	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 22C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾

Page: 32
Run Date: 8 / 18 / 10

Iowa
Summary of Trends in Smoking Indicators by Race/Ethnicity

Summary of Trends in Smoking Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Asian/Pacific Islander	2009	286	10.5	328	5.5	21	*	275	12.0	325	4.3
	2008	237	9.3	305	3.9	18	*	240	13.8	302	9.6
	2007	213	11.3	264	4.9	13	*	172	14.0	257	9.3
	2006	83	*	104	8.7	0	*	0	*	99	*
	2005	312	9.6	248	4.0	18	*	247	21.5	248	17.3
	2004	275	14.2	258	5.4	22	*	170	25.9	258	21.7
	2003	416	11.1	407	4.2	35	*	312	24.0	407	20.1
	2002	396	10.1	381	3.1	22	*	283	26.5	381	17.1
	2001	308	9.7	304	3.6	18	*	225	27.6	304	22.4
	2000	365	12.9	352	5.1	31	*	269	26.4	352	23.0
Multiple Races	2009	193	45.1	220	27.3	65	*	185	22.2	218	9.6
	2008	190	43.2	203	28.1	65	*	188	28.7	198	18.2
	2007	180	45.0	216	25.9	61	*	150	36.0	215	19.5
	2006	62	*	73	*	0	*	0	*	72	*
	2005	0	*	0	*	0	*	0	*	0	*
	2004	0	*	0	*	0	*	0	*	0	*
	2003	0	*	0	*	0	*	0	*	0	*
	2002	0	*	0	*	0	*	0	*	0	*
	2001	0	*	0	*	0	*	0	*	0	*
	2000	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 22C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾

Page: 33
Run Date: 8 / 18 / 10

Iowa
Summary of Trends in Smoking Indicators by Race/Ethnicity

Summary of Trends in Smoking Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
All Other/Unknown	2009	1	*	0	*	0	*	1	*	0	*
	2008	2	*	1	*	0	*	2	*	1	*
	2007	0	*	0	*	0	*	0	*	0	*
	2006	0	*	0	*	0	*	0	*	0	*
	2005	0	*	0	*	0	*	0	*	0	*
	2004	0	*	0	*	0	*	0	*	0	*
	2003	0	*	0	*	0	*	0	*	0	*
	2002	0	*	0	*	0	*	0	*	0	*
	2001	0	*	0	*	0	*	0	*	0	*
2000	0	*	0	*	0	*	0	*	0	*	
Total All Race/Ethnic Groups	2009	15,549	39.8	17,863	23.1	5,069	30.7	15,142	20.8	17,694	8.8
	2008	15,279	40.8	17,175	25.1	5,054	29.8	15,073	24.4	17,058	15.4
	2007	15,104	42.8	18,082	25.6	4,619	30.7	12,455	26.3	17,812	16.8
	2006	4,580	41.0	6,096	25.7	0	*	6	*	6,002	16.5
	2005	17,506	42.1	14,294	27.1	4,827	29.4	14,443	41.9	14,289	36.5
	2004	12,492	41.9	11,435	26.9	3,252	29.1	9,111	41.9	11,432	36.0
	2003	17,451	43.1	16,563	27.1	5,542	29.9	13,966	40.8	16,550	35.8
	2002	17,302	44.2	16,331	27.7	5,615	29.9	13,723	41.7	16,309	34.7
	2001	16,419	46.1	15,026	29.8	5,393	27.3	13,049	44.4	15,021	36.4
	2000	16,981	46.0	15,747	29.6	5,602	27.9	13,443	44.6	15,739	37.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 23C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾

Page: 34
Run Date: 8 / 18 / 10

Iowa

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
White, Not Hispanic	2009	9,842	7.5	8.8	8,767	10.2	7,562	2.9	10,455	30.4
	2008	10,054	7.7	8.4	8,619	10.8	7,559	2.9	10,388	26.7
	2007	10,626	7.7	8.6	7,869	10.5	6,931	2.9	10,931	19.0
	2006	3,526	7.3	8.7	2,499	9.8	2,222	2.4	3,604	21.3
	2005	10,682	7.3	8.6	10,360	12.9	9,007	2.9	10,667	55.4
	2004	8,509	7.6	8.6	8,040	12.9	6,992	2.8	8,490	55.6
	2003	12,556	6.6	9.5	12,073	12.3	10,562	2.6	12,555	54.9
	2002	12,398	7.3	9.4	11,876	12.6	10,343	2.9	12,392	54.5
	2001	11,699	6.7	10.1	11,277	11.7	9,943	3.1	11,672	53.6
	2000	12,344	6.7	10.1	11,885	11.3	10,528	2.7	12,323	52.4
Black, Not Hispanic	2009	966	11.9	4.9	884	13.2	727	6.6	1,041	28.0
	2008	908	11.2	4.7	794	13.6	674	4.9	936	26.5
	2007	995	12.0	4.9	738	13.6	630	4.3	1,027	21.5
	2006	307	9.1	5.2	205	9.8	181	4.4	320	25.3
	2005	987	11.9	4.9	959	16.4	802	5.7	985	47.1
	2004	776	12.9	3.7	718	17.1	593	4.7	779	46.1
	2003	1,079	11.9	4.5	1,040	14.5	886	5.1	1,083	48.6
	2002	1,027	10.1	5.6	974	14.4	830	4.0	1,024	49.7
	2001	960	13.1	4.4	939	19.1	755	5.4	962	44.5
	2000	968	11.3	4.6	938	14.9	792	4.9	970	44.6

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾

Page: 35
Run Date: 8 / 18 / 10

Iowa

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
Hispanic	2009	2,144	6.4	9.2	1,993	9.5	1,728	3.1	2,290	48.3
	2008	2,222	7.4	7.9	1,985	11.6	1,726	2.4	2,306	43.6
	2007	2,390	6.1	9.3	1,916	11.2	1,668	1.7	2,464	38.1
	2006	866	5.5	9.0	694	10.7	607	1.8	892	42.2
	2005	1,866	5.3	8.9	1,826	11.8	1,595	2.4	1,881	78.0
	2004	1,447	6.5	8.8	1,359	11.4	1,197	2.5	1,452	81.0
	2003	1,920	5.8	9.1	1,840	11.8	1,611	2.4	1,934	79.0
	2002	1,908	5.7	10.3	1,830	12.5	1,584	2.2	1,922	78.1
	2001	1,583	4.6	9.3	1,523	11.5	1,341	1.9	1,581	77.3
	2000	1,537	5.9	9.8	1,485	11.0	1,315	2.1	1,538	74.0
American Indian/Alaska Native	2009	66	*	*	59	*	52	*	71	*
	2008	77	*	*	67	*	61	*	80	*
	2007	76	*	*	54	*	49	*	79	*
	2006	21	*	*	15	*	11	*	21	*
	2005	88	*	*	84	*	68	*	88	*
	2004	63	*	*	59	*	49	*	64	*
	2003	84	*	*	79	*	71	*	83	*
	2002	94	*	*	83	*	65	*	94	*
	2001	68	*	*	66	*	55	*	67	*
	2000	72	*	*	67	*	60	*	70	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾

Page: 36
Run Date: 8 / 18 / 10

Iowa

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
Asian/Pacific Islander	2009	259	6.2	2.7	214	8.4	192	2.6	269	44.2
	2008	222	9.0	3.2	184	12.0	156	6.4	235	40.4
	2007	209	10.5	2.9	144	6.9	132	3.0	211	32.2
	2006	70	*	*	42	*	37	*	73	*
	2005	238	10.1	3.8	233	15.0	198	7.1	237	65.8
	2004	246	7.3	5.3	226	14.2	194	2.1	245	67.8
	2003	391	9.0	3.1	373	15.3	315	3.8	393	66.7
	2002	369	8.4	4.3	349	16.0	293	4.4	368	61.4
	2001	294	7.1	3.1	277	15.5	233	2.1	295	59.3
	2000	343	9.3	5.0	333	12.9	288	4.2	344	52.9
Multiple Races	2009	163	8.6	7.4	152	8.6	131	6.1	173	27.7
	2008	161	8.7	5.0	148	10.1	131	2.3	168	26.8
	2007	172	10.5	7.6	127	11.8	111	5.4	177	16.4
	2006	58	*	*	47	*	40	*	58	*
	2005	0	*	*	0	*	0	*	0	*
	2004	0	*	*	0	*	0	*	0	*
	2003	0	*	*	0	*	0	*	0	*
	2002	0	*	*	0	*	0	*	0	*
	2001	0	*	*	0	*	0	*	0	*
	2000	0	*	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2009 Pregnancy Nutrition Surveillance ⁽¹⁾

Page: 37
Run Date: 8 / 18 / 10

Iowa

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
All Other/Unknown	2009	0	*	*	0	*	0	*	0	*
	2008	1	*	*	0	*	0	*	1	*
	2007	0	*	*	0	*	0	*	0	*
	2006	0	*	*	0	*	0	*	0	*
	2005	0	*	*	0	*	0	*	0	*
	2004	0	*	*	0	*	0	*	0	*
	2003	0	*	*	0	*	0	*	0	*
	2002	0	*	*	0	*	0	*	0	*
	2001	0	*	*	0	*	0	*	0	*
	2000	0	*	*	0	*	0	*	0	*
Total All Race/Ethnic Groups	2009	13,440	7.6	8.5	12,069	10.2	10,392	3.2	14,299	33.3
	2008	13,645	7.9	8.0	11,797	11.1	10,307	3.0	14,114	29.7
	2007	14,468	7.8	8.4	10,848	10.8	9,521	2.8	14,889	22.5
	2006	4,848	7.2	8.4	3,502	10.1	3,098	2.5	4,968	25.6
	2005	13,861	7.4	8.3	13,462	13.1	11,670	3.1	13,858	58.1
	2004	11,041	7.8	8.2	10,402	13.0	9,025	2.9	11,030	58.6
	2003	16,030	6.9	9.0	15,405	12.4	13,445	2.7	16,048	57.7
	2002	15,796	7.3	9.2	15,112	12.8	13,115	2.9	15,800	57.2
	2001	14,604	6.9	9.5	14,082	12.3	12,327	3.1	14,577	55.7
	2000	15,264	6.9	9.6	14,708	11.5	12,983	2.8	15,245	54.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

